

Isolation and Identification of Flavonoids from *Arctiumlappa* Stem and Study the Hepatoprotective Effect on Acetaminophen Induced Liver Damage

Zaid Abd Al Salam Alsamarrai¹, Abdulsalam Tawfeeq Alsamaraian², Rafah Razooq Al-Samarrai³

¹Researcher. Department of Chemistry, College of Education, University of Samarra, Samara, Iraq, ²Lect. Dr. Department of Chemistry, College of Education, University of Samarra, Samara, Iraq, ³Prof. Dr. Department of Chemistry, College of Education, University of Samarra, Samara, Iraq

Abstract

The present study investigate the hepatoprotective and antioxidant effect of crude extract of *Arctiumlappa* stem (E-ALS) and isolated total flavonoids (ITF) from the stem, With qualitative and quantitative determination of flavonoids by HPLC. Twenty four local adult male rabbits were used in this study which divided into four groups (6 animal in each group), Control-1 (C1) as control group, Control-2 (C2): treated with 300mg/kg paracetamol administrated for one week, Group 1 (G1): 300mg/kg paracetamol administrated for one week + Orally 250mg/kg of E-ALS administrated daily for 4 weeks, Group 2 (G2): Orally 300mg/kg paracetamol administrated for one week + Orally 50mg/kg of ITF administrated daily for 4 weeks.

The results identified five types of flavonoids for the first time in plant stem (Rutin, Myricetin, Quercetin, Apigenin, Kaempferol) with quantified determination for the concentration of each type of flavonoids in E-ALS and also in ITF which identified the same five flavonoids. The hepatoprotective effect of E-ALS and ITF were observed by monitoring the antioxidant parameters and activity of liver enzymes.

The results obtained from this study showed that the levels of glutathione-GSH, Glutathione peroxidase-GPX and Glutathione-S-Transferase-GST were significantly ($P \leq 0.05$) decreased in C2 treated with paracetamol as comparing with C1, with significant ($P \leq 0.05$) elevation of alanine aminotransferase-ALT activity and non-significant effect on Aspartate aminotransferase-AST, while the level of GSH and GPX were significantly ($P \leq 0.05$) elevated in G1 and G2 treated with E-ALS and ITF respectively as compared with C2, in which the levels of GSH and GPX are less than the levels in C1 and G1. Otherwise the levels of GST, ALT and AST significantly ($P \leq 0.05$) decreased in G1 and G2 as compared with C2. These findings show the hepatoprotective properties of E-ALS and IFT against liver injury induced by paracetamol and also the protective role of anti-oxidative defense system of flavonoids in the two extracts.

Keywords: *Arctiumlappa*, flavonoids, acetaminophen, Glutathione, Glutathione-S-transferase, liver enzymes.

Introduction

The flavonoids are a category of natural substances belonging to the family of polyphenols, they are one of the important types of plant secondary metabolites, widely distributed in foods and Medicinal Plants^(1,2). More than thirty years ago, the research studies focusing on flavonoids from medicinal plant species have increased considerably, because of their versatile benefits for human health⁽³⁾. These includes: antioxidants, anti-inflammatory, anticancer, antibacterial, antiviral, anti-allergic, immune system promoting and also as detoxifying and pro-survival agents^(4,5,6). So that the many researcher use different method of isolation and identification of flavonoids from different medicinal plants and evaluate the pharmacological effect of the isolated flavonoids, such as; Al-Salihi et al, study the Hypolipidemic effect of isolated flavonoids from date palm pollen⁽⁷⁾, while AlSamarraietal⁽⁸⁾, isolated the flavonoids from Bay leaf *Laurusnobilis* L. and also study the hypolipidemic effect.

Arctiumlappa (also known as burdock), one of the important plant in traditional medicine worldwide medicine,

many studies have evaluate the biological activities of the different parts of plant, roots, seeds and leaves^(9,10), including antioxidant activities⁽¹¹⁾, anti-inflammatory⁽¹²⁾, anti-cancer⁽¹³⁾ and anti-hepatotoxicity⁽¹⁴⁾. Many researcher identified many secondary metabolites in different parts of the plant, which include: phenolic compounds, lignans, saponins, tannin, sterols, alkaloids and Flavonoids⁽¹⁵⁻¹⁷⁾, The types of flavonoids which identified in *Arctiumlappaleavesinclude*, luteolin, rutin, quercetin and quercetinrhannoside. On the other hand Rajasekharan^{etal}⁽¹⁸⁾, identified quercetin 3-vicianoside and quercetin 3-O-glucuronide in the root of the plants, with no flavonoids in seeds. The present study aim to isolate and identify the flavonoids from *Arctiumlappastem* and study the protective effect of it against the liver damage induced by *paracetamol*(acetaminophen).

Material and Method

Plant Materials: The dried stems of *Arctiumlappawere* obtain from a local market in Samarra city, Salah Al-Din, Iraq. The stems were separated from other parts of the plant, kept in a dark container until used.

Method

I. Preparation of extracts:

- Crude extract from *Arctiumlappastems E-ALS*: 45g of *Arctiumlappastems powder* was suspend in 180ml normal saline solution.
- Isolation of total flavonoids from *Arctiumlappastems*: This Isolation was done according to (Chen *et al*, method) with some modification⁽¹⁹⁾, the first step before extraction was remove fatty contents, 250g plant stems powder were extracted with 750ml diethyl ether, soxhlet apparatus for 3 hours, dried the defatted plant material at 35 C° in an air oven, the second step extracted flavonoids twice with 500ml (70%) ethanol solution at 90C° for 2h. The solution was filtered and centrifugation at 3000 rpm for 15 min. The solvent was evaporated and the extract was condensed under reduced pressure. The extract was collected, labeled as ITF and stored at 4C° until used.

II. Identification of flavonoids by High-Performance Liquid Chromatography-HPLC: Identification of flavonoids in *Arctiumlappastems* and in isolated flavonoids were carried out according⁽²⁰⁾ method with some modification, In which 500mg of plant

sample(crude, isolated flavonoids extract from stems) were dissolved in 20 ml of hexane to remove fat, 100 ml of 80:20 (methanol: water), the extract was subjected to ultra-sonication at 60% duty cycles for 25 min at 25° c followed by centrifugation at 7,500 rpm for 15min. The clear supernatant of each sample was subjected to charcoal treatment to remove pigments prior to evaporation under vacuum (BuchiRotavapor Re Type). Dried samples were re-suspended in 1.0 ml HPLC grade methanol by vortexing, the mixture were passed through 2,5um disposable filter and stored at 4° C for further analysis, then 20 µl of the sample injected into HPLC system according the optimum condition. Five stander solutions (25µg/ml) were used (Rutin, Myricetin, Querecetin, Apigenin and Kaempferol).

The concentration of identified flavonoids was done according to the following equation:

$$\text{Area of sample} = (\text{Conc. of Flavonoids}(\mu\text{g/ml}) / \text{Area of standard}) \times C \times D$$

C = Conc. of Standard solution

D = Dilution factor

III. Animals: twenty four local adult male rabbits (1200-1550 g weight) were used in this study, Groups of rabbits were housed at room temperature with a lighting schedule of 12 hours light and 12 h dark. Animals had free access to a standard pellet diet. All animals were divided into four groups (6 animal in each group) described as follow:

Control 1(C1): Orally 1ml/kg/day administrated daily dose normal saline only.

Control 2(C2): Orally 1ml/kg/day of *paracetamol* (300mg/kg- normal saline was used as solvent) administrated daily for 7 days.

Group 1(G1): Orally 1ml/kg/day of *paracetamol* (300mg/kg- normal saline was used as solvent) administrated daily for 7 days+Orally 1ml/kg/day of *Arctiumlappastems* extract (250mg/kg- normal saline was used as solvent) administrated daily for 4 weeks.

Group 2(G2): Orally 1ml/kg/day of *paracetamol* (300mg/kg-normal saline was used as solvent) administrated daily for 7 days+Orally 1ml/kg/day of flavonoids isolated from *Arctiumlappastems* (50mg/kg-normal saline was used as solvent) administrated daily for 4 weeks.

IV. Collection of blood samples: After 4 weeks, serum samples were collected by heart puncture from fasting rabbits for 12 hours. Determination of serum levels of Glutathione-GSH^(21,22), Glutathioneperoxidase-GPX⁽²³⁾, Glutathione-Transferase-GST⁽²⁴⁾, alanine aminotransferase-ALT and Aspartate aminotransferase-AST⁽²⁵⁾ by using standard method.

V. Statistical Analysis: Results were analyzed statistically by using (analysis of variance test-ANOVA the statistical program Minitab). Averages were compared to calculations of the characteristics of the application Duncan's Multiple Range Test by probability level $P \leq 0.05$.

Results and Discussion

Part I: Phytochemical study: The HPLC analysis of flavonoids were done firstly by using five standard

flavonoids, Table 1 showed the Retention Times and Area Under Curves for Standard Flavonoids

Table 1: Retention Times and Area Under Curves for Standard Flavonoids

Standard Flavonoids	Retention time (min)	Area under curve μ volt
Rutin	2.473	256193
Myricetin	3.742	261362
Quercetin	4.80	276233
Apigenin	5.877	240672
Kaempferol	6.64	249241

The HPLC analysis of the crude *Arctiumlappa* stem extract showed seven peaks with different Rt (1.987, 2.51, 3.807, 4.883, 5.763, 6.038, 6.823)min [Fig. 1], while the area under curve were (186616, 406614, 387611, 58062, 404152) μ volt [Table 2].




Fig. 1: HPLC analyses of flavonoids in *Arctiumlappa* stems

The types of flavonoids in *Arctiumlappa* stems were identified by comparing the R_t obtained from chromatograms of *crude stems extract* with R_t in chromatogram of standard flavonoids, and then the concentration of the identified flavonoids was done by using the values of the area under curve for stem

extract and for each standard. The results indicate that the crude plants contain 793.56 μ g/g of rutin, 741.33 μ g/g of Myricetin, 866.31 μ g/g of Quercetin, 44.80 μ g/g Apigenin and 810.76 μ g/g Kaempferol with two unknown peaks [Table 2].

Table 2: Retention time, area under curve and concentration of identified flavonoids in *Arctiumlappastems*

Identified compounds	Retention time (min)	Area μ volt	Concentration g/ μ
UnKnown	1.987	186616
Rutin	2.51	406614	793.56
Myricetin	3.807	387510	741.33
Quercetin	4.883	478611	866.31
Apigenin	5.763	21568	44.80
Kaempferol	6.038	58062	810.76
UnKnown	6.823	404150

The HPLC analysis of flavonoids in isolated flavonoids from plant stem showed six peaks with different Rt(2.843, 2.490, 3.828, 4.98, 6.088, 6.852) and area under curve were(121378, 311395, 189935, 397924, 241849, 318015) Table 3.

The result indicate that the types of flavonoids in isolated flavonoids fraction from the plant stem indicate 607.73 μ g/g of rutin, 363.35 μ g/g of Myrecetin, 720.26 μ g/gQuercetin, 502.44 μ g/gApigenin and 637.96 μ g/g Kaempferol, with two unknown peaks [table 3].

Table 3: Retention time, area under curve and concentration of identified flavonoids in isolated flavonoids from *Arctiumlappastems*

Identified compounds	Retention time (min)	Area μ volt	Concentration g/ μ
UnKnown	2.843	121378
Rutin	2.49	311395	607.73
Myricetin	3.828	189935	363.35
Quercetin	4.983	397924	720.26
Apigenin	6.888	241849	502.44
Kaempferol	6.852	318015	637.96

Part II: Biochemical study: The hepatoprotective effect of crude extract and isolated flavonoids from *Arctiumlappastems* as an antioxidant on liver damage induced by acetaminophen were investigate in the present study, the results indicate that the levels of GSH, GPX and GST were significantly ($P \leq 0.05$) decreased in control-2 treated with paracetamol as comparing with control-1 group with significant($P \leq 0.05$) elevation of ALT activity and non-significant effect on AST, while the level of GSH and GPX were significantly($P \leq 0.05$) elevated in G1 and G2 treated with crude stem extract and isolated flavonoids respectivelyas compared with C2, in which the levels of GSH and GPXare less than the levels in C1 and G1. Otherwise the levels of GST,ALT and AST significantly($P \leq 0.05$) decreased in G1 and G2 as compared with C2.

Discussion

Arctiumlappa one of the common medicinal plant in china. The fresh or dried roots, ripe seed and leaf were used medicinally⁽²⁶⁾, with no information related to use of plantstem in traditional medicine, in spite of the *Arctiumlappa* stem maybe eaten stewed or raw as a snack⁽²⁷⁾. Kim *etal*⁽²⁸⁾, indicate that the methanolic extract from leaves and stem of *Arctiumlappa* have anti-inflammatory effect. In recent years, the drug derived from natural sources have been given much attentions more than the chemical drugs, in addition to study the chemical composition and biochemical effect. The *Arctiumlappa* contain lignin especially arctiin (lower concentration than other part of the plant)⁽²⁹⁾, no more information were available about the chemical composition of the *Arctiumlappa* stem or about the

protective effect of it against the liver damage induced by *paracetamol*.

The results of the present study identified five types of flavonoids for the first time in plant stem (Rutin, Myricetin, Quercetin, Apigenin, Kaempferol) with quantified determination for the concentration of each type of flavonoids by HPLC, isolated the stem flavonoids and identified the same five flavonoids and then study the hepatoprotective and antioxidant effect of the crude stem and its isolated flavonoids.

Barrosetal⁽³⁰⁾ provides invaluable insights into the therapeutic efficacy of quercetin in acetaminophen induced toxic liver damage, which suggest that the reduction in serum ALT and AST levels due to the treatment with quercetin, which is useful for prevention of liver damage caused by paracetamol, this results agree with the result of the present study, whereas the treatment with crude extract of stem and isolated flavonoids significantly ($P \leq 0.05$) reduced the activity of AST and ALT in serum of rabbit in G1 and G2, on the other hand Wang *et al*⁽³¹⁾ indicate that treatment with different doses of kaempferol decrease the oxidative stress, lipid peroxidation and increase the antioxidant defense activity, So this findings show the protective effect of kaempferol against liver injury.

The treatment with high dose of paracetamol in C1 cause significant high ($P \leq 0.05$) elevation in serum liver enzymes (AST and ALT) and reduction of antioxidant parameters levels (GSH, GPX, GST), this depression in the level of GSH may be due to the effect of toxic metabolite N-acetyl-p-benzoquinone imine-NAPQI, which produced by hepatic cytochrome p450- CYP450 system which oxidize the excess paracetamol, and the normal detoxification of the toxic metabolite (NAPQI) by the thiol group in GSH cause consume it⁽³²⁾, resulting in accumulation of NAPQI which then binds covalently -SH groups of proteins in hepatocytes forming NAPQI-protein adducts

Conclusion

1. *Arctiumlappastem (E-ALS) and isolated total flavonoids (ITF)* have hepatoprotective properties against liver injury induced by paracetamol.
2. *Arctiumlappastem (E-ALS) and isolated total flavonoids (ITF)* have the protective role of antioxidative defense system.

Ethical Clearance: The Research Ethical

Committee at scientific research by ethical approval of both MOH and MOHSER in Iraq.

Conflict of Interest: Non

Funding: Self-funding

References

1. Fernández SP, Wasowski C, Loscalzo LM, Granger RE, Johnston GAR, Paladini AC, et al. Central nervous system depressant action of flavonoid glycosides. *Eur J Pharmacol.* 2006;539(3):168–76.
2. Burak M, Imen Y. Flavonoids and their antioxidant properties. *Turkiye Klin Tip Bil Derg.* 1999;19:296–304.
3. Ahmed SI, Hayat MQ, Tahir M, Mansoor Q, Ismail M, Keck K, et al. Pharmacologically active flavonoids from the anticancer, antioxidant and antimicrobial extracts of *Cassia angustifolia* Vahl. *BMC Complement Altern Med.* 2016;16(1):460.
4. Kumar S, Pandey AK. Chemistry and biological activities of flavonoids: an overview. *Sci World J.* 2013;2013.
5. Meng X-H, Liu C, Fan R, Zhu L-F, Yang S-X, Zhu H-T, et al. Antioxidative Flavan-3-ol Dimers from the Leaves of *Camellia fangchengensis*. *J Agric Food Chem.* 2018;66(1):247–54.
6. Bjørklund G, Dadar M, Chirumbolo S, Lysiuk R. Flavonoids as detoxifying and pro-survival agents: What's new? *Food Chem Toxicol.* 2017;110:240–50.
7. Al-Salihi FG, Hameed RR. Hypolipidemic effect of date palm pollen and isolated flavonoids in sera of adult male rabbits. *karbala J Pharm Sci.* 2013;(5):34–45.
8. AL-Samarrai OR, Naji NA, Hameed RR. Effect of Bay leaf (*Laurus nobilis* L.) and its isolated (flavonoids and glycosides) on the lipids profile in the local Iraqi female rabbits. *Tikrit J Pure Sci.* 2018;22(6):72–5.
9. Brasileiro BG, Pizziolo VR, Raslan DS, Jamal CM, Silveira D. Antimicrobial and cytotoxic activities screening of some Brazilian medicinal plants used in Governador Valadares district. *Rev Bras Ciências Farm.* 2006;42(2):195–202.
10. Ferracane R, Graziani G, Gallo M, Fogliano V, Ritieni A. Metabolic profile of the bioactive compounds of burdock (*Arctium lappa*) seeds, roots

- and leaves. *J Pharm Biomed Anal.* 2010;51(2):399–404.
11. Liu W, Wang J, Zhang Z, Xu J, Xie Z, Slavin M, et al. In vitro and in vivo antioxidant activity of a fructan from the roots of *Arctium lappa* L. *Int J Biol Macromol.* 2014;65:446–53.
 12. De Almeida ABA, Sanchez-Hidalgo M, Martín AR, Luiz-Ferreira A, Trigo JR, Vilegas W, et al. Anti-inflammatory intestinal activity of *Arctium lappa* L. (Asteraceae) in TNBS colitis model. *J Ethnopharmacol.* 2013;146(1):300–10.
 13. Sun Q, Liu K, Shen X, Jin W, Jiang L, Sheikh MS, et al. Lappaol F, a novel anticancer agent isolated from plant *Arctium Lappa* L. *Mol Cancer Ther.* 2014;13(1):49–59.
 14. F de Souza Predes F, da Silva Diamante MA, Foglio MA, Camargo CA, Aoyama H, Miranda SC, et al. Hepatoprotective effect of *Arctium lappa* root extract on cadmium toxicity in adult Wistar rats. *Biol Trace Elem Res.* 2014;160(2):250–7.
 15. Wang HY, Yang JS. Studies on the chemical constituents of *Arctium lappa* L. *Yao xue xue bao= Acta Pharm Sin.* 1993;28(12):911–17.
 16. Park SY, Hong SS, Han XH, Hwang JS, Lee D, Ro JS, et al. Lignans from *Arctium lappa* and their inhibition of LPS-induced nitric oxide production. *Chem Pharm Bull.* 2007;55(1):150–52.
 17. Al-Snafi AE. The Pharmacological importance and chemical constituents of *Arctium Lappa*. A review. *Int J Pharm Res Sch.* 2014;3(1–1):663–70.
 18. Rajasekharan SK, Ramesh S, Bakkiyaraj D, Elangomathavan R, Kamalanathan C. Burdock root extracts limit quorum-sensing-controlled phenotypes and biofilm architecture in major urinary tract pathogens. *Urolithiasis.* 2015;43(1):29–40.
 19. Chen JJ, Li XR, Fang X. Purification of total flavones from *Morus alba* L. by macroporous adsorbents and kinetic model for the process. *Zhejiang da xue xue bao Yi xue ban= J Zhejiang Univ Med Sci.* 2006;35(2):219–23.
 20. Rodríguez-Delgado MA, Malovana S, Perez JP, Borges T, Montelongo FJG. Separation of phenolic compounds by high-performance liquid chromatography with absorbance and fluorimetric detection. *J Chromatogr A.* 2001;912(2):249–57.
 21. Burtis CA, Ashwood ER. *Tietz textbook of clinical chemistry.* Philadelphia. 1999;1999:1654–50.
 22. Seadlak J, Lindsay RH. *Analytical Biochemistry.* 192, Cited by Al-Zamyle, OM, Al-Nimer MS, Al-Muslih RK (2001). Detection the level of peroxynitrite and related with antioxidant satus in the serum of patients with acute myocardial infraction. *Nation J Chem.* 1968;4:625–37.
 23. Green MJ, Hill HAO. [1] *Chemistry of dioxygen.* In: *Method in enzymology.* Elsevier; 1984. p. 3–22.
 24. Habig WH, Pabst MJ, Jakoby WB. Glutathione S-transferases the first enzymatic step in mercapturic acid formation. *J Biol Chem.* 1974;249(22):7130–39.
 25. Reitman S, Frankel S. A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *Am J Clin Pathol.* 1957;28(1):56–63.
 26. Fleming, T. (2000). *PDR for herbal medicines.* PDR for herbal medicines. (Ed. 2) 128-129.
 27. Tardío J, Pascual H, Morales R. Wild food plants traditionally used in the province of Madrid, Central Spain. *Econ Bot.* 2005;59(2):122-136.
 28. Kim Y-K, Koppula S, Shim D-W, In E-J, Kwak S-B, Kim M-K, et al. Inhibitory effect and mechanism of *arctium lappa* extract on nlrp3 inflammasome activation. *Evidence-Based Complement Altern Med.* 2018;2018.
 29. Gao Q, Yang M, Zuo Z. Overview of the anti-inflammatory effects, pharmacokinetic properties and clinical efficacies of arctigenin and arctiin from *Arctium lappa* L. *Acta Pharmacol Sin.* 2018;39(5):787–801.
 30. Barros PP, Silva GH da, Gonçalves GMS, Oliveira JC, Pagnan LG, Arco-e-Flexa L. Hepatoprotective effect of quercetin pretreatment against paracetamol-induced liver damage and partial hepatectomy in rats. *Brazilian Arch Biol Technol.* 2017;60:1-10.
 31. Wang M, Sun J, Jiang Z, Xie W, Zhang X. Hepatoprotective effect of kaempferol against alcoholic liver injury in mice. *Am J Chin Med.* 2015;43(02):241–54.
 32. Yanpallewar SU, Sen S, Tapas S, Kumar M, Raju SS, Acharya SB. Effect of *Azadirachta indica* on paracetamol-induced hepatic damage in albino rats. *Phytomedicine.* 2003;10(5):391–96.