

Effect of Carcinogenic Substance (7,12 Dimethylbenz [a] Anthracene (DMBA)) on Tissue, Hematology Character and Enzyme Activity in Rat

Abdulridha Mohammed Al-Asady¹, Ntisar Kadhum Ghaleb², Abeer Muhammedali Jasim Alnasrawi³, Thoalffakar A. ALhamed⁴

¹Assist.Lecturer, College of Nursing, University of Warith Al-anbiya 'a , Iraq, ²Assist.Lecturer, Department of Biology, College of Science, University of Kerbala, Iraq, ³Assist.Lecturer, Department of Clinical Laboratories, College of Applied, Medical Science, University of Kerbala, Iraq, ⁴ College of Nursing, University of Warith Al-Anbiya 'a, Iraq

Abstract

This study aimed to investigate the effects of 7,12 dimethylbenz [a] anthracene (DMBA) on some blood parameters and hepatic histopathology in rats and antioxidant enzyme. twenty female Wistar albino rats, weighing 180–200 g, were randomly divided into two group. DMBA group (positive group) who received 20 mg DMBA/kg body weight/ (single Douse). Control group (negative group) don't received any thing. The animals in the groups were sacrificed at the end of the 90 days: The histological structure of the liver tissues in the control group was normal. the liver exhibited hydropic degeneration and coagulation necrosis in hepatocytes, severe dilation in the sinusoids, congestion in the central and portal regions , DMBA groups, was degenerative and necrotic changes were detected. MDA levels increased in the DMBA group , compared to the control group, Antioxidant activity CAT, SOD and GPX, the results showed increased enzyme activity in positive control compared with negative groups, ,the hematological parameter was increase in WBC, PLT, GRA, and LYM and decrease in RBC value .

Keywords: 7,12-dimethylbenz[a]anthracene (DMBA); Blood parameters; Antioxidant enzyme ; MDA ; Rat.

Introduction

Cancer, also termed as malignant neoplasm, is a type of diseases where in a group of cells show abnormal proliferation, invasion and sometimes metastasis 1 . Cancer begins when cells in a part of the body start to grow out of control 2 . There are more than 100 types of cancer have been identified 3 . The substances that promote cancers are called carcinogens, and agents that have the ability to change DNA in ways that are inherited by daughter cells are called mutagens. Most cancers are related to environmental, lifestyle, or behavioral exposures 4 The term “environmental”, as used by cancer researchers, refers to everything outside the body that interacts with humans 5 . Other chemicals, like benzene, ketones, vinyl chloride, ethylene bromide, and dichloro-diphenyl-trichloroethane (DDT), are known carcinogens 6 . The accumulation of multiple factors, carcinogens and altered genes, transform a normal cell into a cancerous one. Carcinogens that have

the ability to promote cancer have various sources 7 .The factors responsible for cancer development are classified as exogenous and endogenous 8 . The first group includes nutritional habits (food preservation and preparation), socio-economic status, lifestyle, physical agents (ionising and non-ionising radiation), chemical compounds (natural and synthetic) and biological agents (Helicobacter pylori, Epstein Barr virus, human T lymphotropic viruses I and II, human papilloma virus and the hepatitis B virus 9.

Meterial and Method

Experiential animals and Tumor induction in rats:

Twenty female of Wistar albino rats and aged between (7-8) weeks were weighing between 150–250 g used in this study. All they were kept in ventilated cages, with temperature of $25\pm 2^{\circ}\text{C}$. A 12:12 h light:dark cycle

is also regulated for these animals. Balanced rodent food pellet and water is provided. The rats were randomly assigned to 2 groups of 10 rats each.

The groups were as follow:

Group 1: Received a single dose of DMBA (positive control) 20 mg DMBA/kg body weight/ (single dose)

Group 2: Did not received any treatment (negative control).

DMBA administration: Mammary tumors were induced by 7, 12-

Dimethylbenz (a) anthracene (DMBA) ¹⁰ . A single dose of DMBA dissolved in corn oil was given by oral gavage to two groups using the syringe and needles. DMBA was purchased from Sigma Aldrich and dissolved in corn oil. The concentration of the solution was 20 mg DMBA per 1 ml corn oil for each rat ¹⁰ .

Preparation of tissue sample


The liver was exteriorized and excised. All specimens were immediately fixed in 10% formaldehyde solution. After fixation they were processed in usual manner, and embedded in paraffin for subsequent histopathological examination for liver. A scoring system (of no abnormalities, mild, moderate, severe) was used to classify the liver changes according to the severity of the damage and extent of histological changes. The histological sections were evaluated by a pathologist without prior knowledge of the treatment given to the animals ¹¹ .

Blood parameters were determined in whole blood by the using rat mode of veterinary practice with a blood cell counter (Abocus Junior Vet-5, Austria). Measurements of biochemical parameters were made with a Modular PP autoanalyzer (Mindray BS800, China).


Results

Histopathological findings: The control group showed normal histological structure in the liver tissue (Figure 1A). In the DMBA group, the livers exhibited dilatation of the sinusoids, cholangiohepatitis in the portal region, and congestion in the sinusoidal and portal regions (Figure 1B). Hematological parameters also change in negative group compared of positive group

Table 1 , in positive group : white blood cell (WBC) , LYM, MON ,GRA and PLT while other parameter such as RBC was decrease. The state of free radicals and antioxidants is given in Table 2. As seen in Table 2, the levels of MDA increased in liver groups compared to the control group, also SOD, GSH-Px, CAT, and GSH



values decreased.



A

B

Figure1; A: DMBA group showing colangiohepatitis in portal region, mild dilation in sinusoids, hyperemia, and hydropic degeneration in hepatocytes;

B: Control group showing liver tissue with a normal histological structure

Table 1. Some hematological parameters in the

groups (values are mean n=4)

No.	Parameter	Negative control	Positive control
1	WBC (103 /mm ³)	8.2	15.1
2	RBC(103 /mm ³)	8.8	1.4
3	LYM %	67%	78.5 %
4	MON%	1	0.9
5	GRA %	18.8 %	94 %
6	MCV(fl)	56.6	58.7
7	MCH(Pg)	16	21
8	MCHC(g/dl)	32.3	32.8
9	PLT(105 /mm ³)	338	737

Table 2. Some biochemical parameters in the groups (values are mean n=4)

Treatments	Conc. of GPx(IU/mg)	Conc. Of SOD (IU/mg)	Conc. of CAT (IU/mg)	Conc. of MDA (nmol/mg)
Negative group	127	458	122	134
Positive group	312.3	567	187	267

Discussion

7, 12-dimethylbenz (α) anthracene (DMBA) is a well known carcinogen and immunosuppressor used in rodent models to study cancer ¹². DMBA is reported to induce mutations by making DNA adducts ^(13, 14). Although, it is a well known skin carcinogen, yet many researchers have reported the deleterious effect of DMBA in liver ^(15, 16). Liver is the primary site of metabolism and is often prone to damage by xenobiotics. Evidently, liver cancer is the second most common cause of cancer deaths worldwide ¹⁷. several haematological and haematochemical parameters were changed when treated with DMBA and found that DMBA induced hepatocellular carcinoma. Experimental studies showed that DMBA-induced skin, oral, mammary and ovarian tumors ¹⁸. The carcinogenic and mutagenic effect of DMBA requires its metabolic activation

by mixed function oxidases. The hydroxylation of DMBA at 7-methyl group is a crucial step towards its carcinogenesis ¹⁹. Further metabolism of DMBA leads to formation of a wide range of metabolites with varying toxicity. Among these, trans-3,4-dihydrodiol-1,2-epoxide is the carcinogenic product of DMBA ^(20,21). The metabolic products of DMBA, when present inside body, hampers ROS-antioxidant balance by overproduction of free radicals and the body in turn reacts by modulating activities of antioxidant enzymes to curb the damaging effects of an increased ROS ²². Hematological and biochemical parameters may be affected by a variety of factors such as race, age, gender, pregnancy, lactation, muscular activity, region, season, environmental heat, maintenance, and nutrition. In the present study, the effects on blood parameters and hepatic, histopathology of fluoride. Oxidative products derived from mutagen metabolism, such as DMBA, might impair vital cellular function by damaging proteins and lipid membranes. Consequently, these changes induced by the chemical carcinogen 7,12-dimethylbenz[a]anthracene, have been reported to be leukemia, and the development of anemia ²³. Reactive oxygen species (ROS) are important as pathological agents for many diseases. Increased oxygen radical production and lipid peroxidation are associated with the pathogenesis of many diseases and the toxic effects of a wide range of compounds increase in total leukocyte count (WBC), eosinophil, neutrophil and monocyte values for rats exposed to DMBA was also reported by ²⁴. Hematological and biochemical parameters may be affected by a variety of factors such as race, age, gender, pregnancy, lactation, muscular activity, region, season, environmental heat, maintenance, and nutrition. a significant increase in the GPX, CAT, SOD level in rat treated with DMBA only was reported by ²⁵. On other hand ²⁶ find MDA content increase significantly with rat treated with DMBA only.

Financial Disclosure: There is no financial disclosure.

Conflict of Interest: None to declare.

Ethical Clearance: All experimental protocols were approved under the College of Nursing, University of Warith Al-anbiya'a, Iraq and all experiments were carried out in accordance with approved guidelines.

References

1. Samudrala PK, Augustine BB, Kasala ER, Bodduluru LN, Barua C, Lahkar M. Evaluation

- of antitumor activity and antioxidant status of *Alternanthera brasiliana* against Ehrlich ascites carcinoma in Swiss albino mice. *Pharm Rese*, 2015; 7(1).
2. Jones S, Johnson K. Women's awareness of cancer symptoms: a review of the literature *Women's Health* 2012; 8(5):579–591
 3. Qi J, Yu S, Zhang F. Reference Gene Selection for Real-Time Quantitative Polymerase Chain Reaction of mRNA Transcript Levels in Chinese Cabbage (*Brassica rapa* L. ssp. *pekinensis*) 2010;28: 597–604
 4. Aggarwal BB, Danda D, Gupta S, Gehlot P.. Models for prevention and treatment of cancer: Problems vs promises. *Biochem Pharmacol*. 2009;78:1083–1094.
 5. National Cancer Institute, Division of Cancer Control and Population Sciences, Applied Research Program, Health Services and Economics Branch., 2013.
 6. Stokstad E. "Species conservation. Can the bald eagle still soar after it is delisted?". *Science*. 2007;316 (5832): 1689–90.
 7. Tomasetti C, Li L, Vogelstein B. "Stem cell divisions, somatic mutations, cancer etiology, and cancer prevention". *Science*. 2017;355 (6331): 1330–1334.
 8. GUTIÉRREZ JB, SALSAMENDI AL. Fundamentos de ciencia toxicológica. Diaz de Santos, Madrid, 2001;155-177.
 9. LUTZWK. Differences in individual susceptibility to toxic effects of chemicals determine the dose-response relationship and consequences of setting exposure standards. *Toxicol Lett* 2002;126: 155–158
 10. Barros AC, Muranaka ENK, Mori LJ, Pelizon CHT, Iriya K, Giocondo G.. Induction of experimental mammary carcinogenesis in rats with 7,12-dimethylbenz (a)anthracene. *Revista do Hospital das Clínicas*. 2004;59 (5):257–61.
 11. Najah RH, Bassim M, Ahmad A. Retard the Progression of Atherosclerosis via Down Regulation of the Inflammatory and Oxidative Pathways. *J Clin Exp Cardiol* 2013;4:6
 12. Chatterjee M, Janarthan M, Manivannan R, Rana A, Chatterjee M Combinatorial effect of fish oil (Maxepa) and 1alpha,25-dihydroxyvitamin D(3) in the chemoprevention of DMBA-induced mammary carcinogenesis in rats. *Chem Biol Interact* 2010;188: 102–110
 13. Morse MA, Baird WM, Carlson GP, Effects of sex and age on DMBA: DNA binding in epidermis of SENCAR mice following topical administration of dimethylbenz[*a*]anthracene. *Cancer Lett* 1987;37: 25–31.
 14. Szafer H, Krajika-Kuzniak V, Ignatowicz E, Adamska T, Markowski J, et al. The effect of cloudy apple juice on hepatic and mammary gland phase I and II enzymes induced by DMBA in female Sprague-Dawley rats. *Drug Chem Toxicol* 2014;37: 472–479.
 15. Singh H, Bedi PS, Singh B Hepatoprotective activity of turmeric and garlic against 7-12, dimethylbenzanthracene induced liver damage in wistar albino rats. *European J Med Plant* 2011;1: 162–170.
 16. Kaur R, Arora S, Interactions of betulinic acid with xenobiotic metabolizing and antioxidative enzymes in DMBA treated Sprague Dawley Female Rats. *Free Rad Biol Med* 2013;65: 131–142
 17. Kumar R, Kaur R, Singh AP, Arora S, Diminution of Hepatic Response to 7, 12-dimethylbenz(a) anthracene by ethyl acetate fraction of *Acacia catechu* Willd. through modulation of xenobiotic and anti-oxidative enzymes in rats. *PLoS ONE* 2014;9: e90083.
 18. Suzuki JS, Nishimura B, Zhang Y. Metallothionein deficiency enhances skin carcinogenesis induced by 7,12-dimethylbenz [a] anthracene and 12-O-tetradecanoylphorbol-13-acetate in Metallothionein-null mice. *Carcinogenesis*, 2003;24: 1123-1132.
 19. Wong LK, Dru J, Lin LS, Knapp J. Metabolism of 7,12-dimethylbenz[a]anthracene by *Cunninghamella elegans*. *Appl Environ Microbiol* 1983;46: 1239–1242.
 20. Huberman E, Chou MW, Yang SK, Identification of 7,12-dimethylbenz[a]anthracene metabolites that lead to mutagenesis in mammalian cells. *Proc Natl Acad Sci USA* 1979;76: 862–866.
 21. Ahmed HA, Manna F, Estefan SF. Modulatory effect of the red sea soft coral extracts on hepatotoxicity induced by carcinogenic agents in rat model. *J Egypt Soc Toxicol* 2006;35: 97–107.
 22. Desai GV, Casciano D, Feuers JR, Aidoo A. Activity profile of glutathione-dependant

- enzyme and respiratory chain complexes in rats supplemented with antioxidants and treated with carcinogens. *Arch. Biochem. Biophys.*, 2001;394, 255-264.
23. Fl-Mofly MM., Abdelmeguid NE , Sadek IA , Essawy AE , Abdel Alem EA.. Induction of leukaemia in chloramphenicol-treated toads East Mediterr. Health J., 2000;6:1026-1034.
24. Batcioglu K , Uyumlu AB , Satilmis B , Yildirim B . Oxidative Stress in the in vivo DMBA Rat Model of Breast Cancer: Suppression by a Voltage-gated Sodium Channel Inhibitor (RS100642), *Basic & Clinical Pharmacology & Toxicology*, 2012;111:137–141.
25. Arroyo-Acevedo , Chávez-Asmat RJ , Anampaguzmán AR , Ráez-González J. Protective Effect of Piper aduncum Capsule on DMBA-induced Breast Cancer in Rats. *Breast Cancer: Basic and Clinical Res.* 2015;9: 41–48 .