

Comparison of Stability And Sternum Healing Rate in Clinical and Ultrasonography (USG) between Stainless Steel Wire and Polydioxanone Yarn in Children Heart Surgery Operation

Prima Kharisma Hayuningrat¹, Heroe Soebroto¹, Paulus Rahardjo²

¹Thoracic Surgery, Cardiovascular and Vascular, ² Department of Radiology, Faculty of Medicine - Dr. Soetomo General Hospital, Universitas Airlangga, Surabaya 60285, Indonesia

Abstract

Background: The closing of the sternum bone is often used today by using stainless steel wire material and Polydioxanone yarn continuous suture. To objectives to be achieved for sternal healing after heart surgery without complications resulting from the failure of sternal healing. The bone healing process itself is influenced by mechanical stress and movement.

Objectives: To compare the effects of sternum closure techniques on surgical patients The heart of the child uses Polydioxanone and Stainless Steel Wire threads against clinical stability and rate of healing which are evaluated clinically and Ultrasonography.

Methods: Performed sternal closure of pediatric patients after cardiac surgery with sternal wire (n = 8) and PDS (n = 8). Performed sternal pain and stability evaluation with the physical examination. Further sternum ultrasonography und was performed to assess displacement, gap and callus picture. Evaluations were performed at weeks 6, 9 and 12 postoperatively.

Results: Week 6 and 9 degrees of pain were higher in sternal wire compared with polydioxanone (p = 0.03 and p = 0.01). The 12th week of sternal wire and polydioxanone did not find any difference in pain (p = 1,000). Week 6, 9 and 12 there was no clinical stability difference between wire and PDS (p = 0.143, p = 0.264, p = 0.063). 9th, 9th and 12th Sternum ultrasonography of examination appears to be displacement in polydioxanone (p = 0.025, p = 0.009, p = 0.009). The gap increased significantly from 6th to 9th weeks in the polydioxanone group, while the addition at week 9 to 12 was statistically insignificant but it appears that polydioxanone had a wider gap addition range than the sternal wire. Week 9 and 12 callus were seen more often in sternal wire patients but not significant (P = 0.602, p = 0.333)

Conclusion: Clinically, sternal steal wire stability is proportional to polydioxanone. Radiologically, the stability of sternal wire is better than polydioxanone. The rate of sternal cure in polydioxanone is proportional to the sternal wire.

Keywords: *Sternal wire, Polydioxanone, Sternum ultrasonography, Paediatrics*

Corresponding Author:

Heroe Soebroto

Thoracic Surgery, Cardiovascular and Vascular,
Faculty of Medicine - Dr. Soetomo General Hospital,
Universitas Airlangga, Jalan Mayjen Prof Dr Moestopo
47, Surabaya Indonesia 60285, Phone: +62315020251
E-mail : unairsubroto@gmail.com

Introduction

Sternotomy was first introduced in 1997 by using experimental goats and human cadaver which then succeeded, then in 1957, performed sternotomy action on heart surgery patients^{1,2}. In the USA, more than 750.000 median sternotomy procedures are performed

annually for cardiac surgery³

Today, closure of the sternum bone that is often used by using stainless steel wire material (SS). The commonly used closure techniques are figure-of-8 and simple interrupted techniques. Biomechanical studies on cadavers and animals show the efficacy of several sternal closure techniques with different results⁴⁻⁶. Currently, the use of synthetic polydioxanone absorbable suture after sternotomy becomes a routine procedure performed in many heart surgery centers, especially in cases of heart surgery in pediatric patients. Several studies have shown the use of polydioxanone (PDS) is effective in preventing sternal instability, assisting wound healing and is more suitable for pediatric patients^{7,8}. In addition, the PDS handling is easier to use, lowering the risk of bleeding at the stitching site and reducing postoperative pain⁹⁻¹¹.

The use of PDS in cardiovascular surgery was first performed on the sternum closure procedure for various heart operations such as correction of coarctation of the aorta, total correction Pulmonary vein drainage anomaly, arterial switches in Transposition of Great Arteries and systemic pulmonary shunts. Postoperative results obtained polydioxanone does not cause inflammatory and mediastinitis reactions, at least granulation tissue and is well-absorbed within the span of 2-6 months. The strength of fixation on the sternum is influenced by the type of fixation technique, the amount, strength and thickness of the wire used and the strength of the sternum itself^{12,13}.

The main goal to be achieved from the sternum closure technique is the achievement of sternal healing after heart surgery without any complications due to sternal healing failure. There are two important factors in bone healing process including mechanical stress and movement which can affect sternum technique stability in callus formation process, the formation or gap between the two sides of the split sternum and the fixation of movement between the sternal fragments^{14,15}.

The routine procedure performed in Dr. Soetomo General Hospital in sternal closure after childhood heart surgery is the most frequent use of continuous suture Polydioxanone yarn while the SSW of the figure of eight is rarely used. Evaluations to assess sternal healing include clinical evaluation of pain scale and physical examination scale assessing sternal stability using the scale¹⁶. This evaluation was used in several studies of

sternal stability assessment after adult heart surgery¹⁷. But there has not been much study in patients after child's heart surgery.

Based on, the study literature comparing the two uses of this material to a child's heart surgical patient as well as evaluating the ratio of its cure. Until now, no research results have been found, therefore, this study intends to compare the effect of sternum closure technique on the patient's heart surgical patients using Polydioxanone and Stainless Steel Wire threads on clinical stability and rate of healing which is evaluated clinically and Ultrasonography.

Method

This study used a double-blinded randomized clinical trial design in patients with cardiac surgery performed by the sternal closure. The study was conducted in the operating room of cardiac surgery, radiology ultrasonography examination room and polyclinic surgery of Toraks, Kardiak, and Vascular Dr. Soetomo General Hospital, Surabaya. The study was conducted from 2016 to 2017. The inclusion criteria included pediatric patients undergoing open and closed heart surgery with a 1-10-year-old sternotomy approach with 10-20 kg. Redo surgery, re-surgery due to postoperative hemorrhage and the patient dies or did not control during the observation period of the study was excluded.

The research procedure, covering the selection phase of the patient according to the inclusion criteria, then the research tool needed sternum closure technique using two types of material include Stainless steel wire and polydioxanone synthetic yarn. Furthermore, in the treatment stage, all patients undergoing standardized cardiac surgery procedures adhere to diagnose and prevailing protocols. The suturing process will use at least 3 SS wire fixations with the figure of -8 and 6 continuous stitching Polydioxanone for each sternum. Furthermore, the observational stage of the study, the patient was evaluated postoperatively with evaluation at the cardiac surgical polyclinic at 9th, 12th week. Sternal ultrasound examination with the transversal projection at three checkpoints (upper sternum, mid sternum, and lower sternum). Statistical data processing by using Mann Whitney Test for the non-parametric test with SPSS v23 program (SPSS, Inc., Chicago, IL.).

Results

The first group of sternal bone was fixed with sternal wire with the stitch of the figure of eight and the second group was fixed with PDS with simple continuous stitching. Patients were followed for postoperative physical and radiological examination at weeks 6, 9 and 12. The sternal closure images with PDS and sternal wire bias are seen in Figures 1 and 2.

The results of stability were obtained in the 6th-week evaluation results in patients using sternal wire, showed a higher degree of pain compared with PDS ($p = 0.03$) with sternal wire group pain characteristics in mild to moderate to severe degrees. While in the PDS group all patients felt a mild degree of pain. Evaluation at week 9 of patients using sternal wire still showed the higher degree of pain than PDS ($p = 0.01$) but with lower pain level characteristics. All patients with sternal wire experience mild pain up to moderate pain level. In the 12th week, both sternal wire and PDS were not found significant differences in pain ($p = 1.000$).

At the 6th-week ultrasound examination, the use of PDS showed a significant displacement compared to the use of sternal wire ($p = 0.025$). At 9th week ultrasound evaluation still showed significant displacement compared to sternal wire ($p = 0.009$) and at week 12 evaluation PDS still showed a significant displacement compared to sternal wire ($p = 0.009$) but no new displacement at other patients.

The results In the sixth week USG examination found a significant gap difference which obtained a wider

gap in PDS in Manubrium ($p = 0.021$) while at mid and lowers there was no significant gap difference ($p = 0.094$ and $p = 0.728$). On the ultrasound examination also has not obtained any callus picture either in the PDS group or sternal wire.

In the 9th-week gap assessment, there was a significant difference with the wider gap of PDS in Manubrium sternum ($p = 0.005$) and on the sternum Corp ($p = 0.001$) while on the lower sternum there was no significant gap difference ($p = 0.069$). In the 12th week, there was a significant difference with the wider gap in the PDS group, i.e., in the Manubrium sternum ($p = 0.001$), the sternum Corp ($p = 0.002$) and the lower sternum ($p = 0.029$).

When compared to the number of gap additions at the 6th, 9th, and 12th-week checks, the gap size increased significantly from week 6 to 9 in the PDS group compared to the sternal wire group. The addition of a gap occurs at all points of the sternal examination. The addition of the gap at the 9th to the 12th week was not statistically significant but it appears that PDS has a wider gap addition range than the sternal wire.

In this 9th week, we have got callus picture on PDS group and sternal wire. The number of patients seen callus in the sternal wire group was more than the PDS, but not statistically significant compared to the PDS group ($P = 0.602$). In the 12th week of ultrasonography results, there was a callus profile of both the PDS and the sternal wire with more callus numbers obtained in sternal wire patients but not statistically significant ($p = 0.333$).

Table 1.Characteristics of the patients

	Sternal wire		PDS	
	Male	Female	Male	Female
Sex	6	2	5	3
Weight (kg) 10 – 20	Mean 12.43 ± 3.150		Mean 12.13 ± 3.064	
Age(y/o) 0 – 12	8.90 ± 1.560		8.60 ± 1.320	

Table 2. Procedure operation

Procedure	Frequency	Percentase
ASD Closure	4	25.0
BCPS	2	12.5
BT Shunt	2	12.5
<i>corpus alienum</i>	1	6.3
TOFTotal Correction	3	18.8
VSD Closure	4	25.0
Total	16	100.0

Table 3. Stability check results

Variables (sternal wire vs PDS) Sternal wire		Week-6		Week-9		Week-12	
		PDS	Sternal wire	PDS	Sternal wire	PDS	
Pain	No				4	8	8
	Mild	2	8	5	4		
	Mod	5		3			
	Sev	1					
	p-price	0.03		0.01		1.00	
Stability	Stable	8	6	7	5	8	5
	Minimal		2	1	3		3
	Partial						
	Complete						
	Harga p	0.143		0.264		0.063	
Displacement	Symmetrical	7	4	7	3	7	3
	Displace	1	4	1	5		5
	p-price	0.025		0.009		0.009	

Table 4. Healing rate results

Variable (sternal wire vs PDS) Sternal wire		Week-6		Week-9		Week-12	
		PDS	Sternal wire	PDS	Sternal wire	PDS	
Callus	Yes			3	2	5	3
	No	8	8	5	6	3	5
	p-price	1.00		0.602		0.333	
Manubrium (mm)		1.31 ± 0.125	1.47 ± 0.12	1.35 ± 0.16	1.75 ± 0.25	1.38 ± 0.20	1.92 ± 0.18
P		0.021		0.005		0.001	
Corpus (mm)		1.13 ± 0.08	1.21 ± 0.09	1.14 ± 0.07	1.38 ± 0.13	1.18 ± 0.06	1.53 ± 0.36
P		0.094		0.001		0.019	
Lower sternum (mm)		1.16 ± 0.05	1.16 ± 0.10	1.16 ± 0.05	1.22 ± 0.07	1.18 ± 0.06	1.32 ± 0.15
P		0.728		0.105		0.029	

Table 5. Average of gap increase

Variable (sternal wire vs PDS)	Week 6 to 9			Week 9 to 12		
	Sternal wire	PDS	p	Sternal wire	PDS	p
Manubrium (mm)	0.04 ± 0.05	0.275±0.183	.004	-0.125 ± 0.125	0.175 ± 0.205	.105
Corpus (mm)	0.00 ± 0.00	0.175 ± 0.167	.004	0.05 ± 0.075	0.150 ± 0.340	.862
Lower sternum (mm)	0.00 ± 0.00	0.062 ± 0.074	.027	0.025 ± 0.07	0.01 ± 0.015	.239

Discussion

The stability and healing of the sternum in this study was stability and clinical and radiological healing evaluation at 6th week, 9th week and 12th week. The sternal stability was evaluated based on the degree of pain on clinical examination as well as the physical examination to assess sternal stability and then evaluate the presence or absence of displacement by examining

ultrasonography. While the healing of the sternum was assessed through ultrasonography examination based on the evaluation of the gap on the sternum as well as the callus picture on the sternal fragment.

The stability and healing process of the sternum is influenced by the combination of daily physiological movements such as breathing, coughing and moving that create a force on the sternum. The force was the

result of a combination of transverse shear, longitudinal shear, and lateral distraction movements. In this study at weeks 6, 9 and 12 seen in ultrasound there was a significant displacement in the PDS group. This study was in accordance with the study wherein the Sternal wire group with strong material properties, proved to have a much better rigidity¹⁸ compared to PDS and this was indicated by a stable symmetry of the sternum.

At week 6 there was a wider gap in the Manubrium and Corpus Sternum groups of PDS than in Sternal wire. The gap differences remained significantly larger at the 9th and 12th weeks. This difference is due to the fact that the PDS uses absorbable material which will certainly be degraded over a period of time. Based on the PDS profile of ethicon® the power of the PDS at week 2 will decrease to 70% and will decrease gradually to 25% at 6th week¹⁹.

At the end of the 12th week of study, the PDS gap compared to the sternal wire was 19.25 ± 1.83 vs 13.75 ± 1.58 . This shows the evaluation of PDS usage by ultrasonography is seen widening and addition of bigger gap than sternal wire. Although the use of Sternal wire proves to be more confined in fixing the Sternum, PDS threads were also able to maintain the gap of Sternum with a distance below 2 mm can trigger the running of the healing process (fibrous tissue healing).

Conclusion

Sternal wire causes higher pain than PDS. Clinically, sternal wire stability was proportional to PDS. Radiologically, the stability of sternal wire was better than PDS. PDS produces a wider sternal gap than a sternal wire. The rate of sternal healing in the PDS was proportional to the sternal wire.

Conflict of Interest: There is no conflict of interest

Source of Funding : This study is self-funded

Ethical Clearance: This study was approved by Ethical Commission of Health Research Faculty of Medicine University of Airlangga.

References

1. Aref R. The role of microcirculation in the prevention and treatment of post-sternotomy mediastinitis. 2018;
2. Julian OC, Lopez-Belio M, Dye WS, Javid H, Grove WJ. The median sternal incision in intracardiac surgery with extracorporeal circulation: a general evaluation of its use in heart surgery. *Surgery*. 1957;42(4):753–61.
3. Bek EL, Yun KL, Kochamba GS, Pfeffer TA. Effective median sternotomy closure in high-risk open heart patients. *Ann Thorac Surg*. 2010;89(4):1317–8.
4. Casha AR, Yang L, Kay PH, Saleh M, Cooper GJ. A biomechanical study of median sternotomy closure techniques. *Eur J Cardio-thoracic Surg*. 1999;15(3):365–9.
5. Glennie S, Shepherd DET, Jutley RS. Strength of wired sternotomy closures: effect of number of wire twists. *Interact Cardiovasc Thorac Surg*. 2003;2(1):3–5.
6. Boiselle PM, Mansilla A V, Fisher MS, McLoud TC. Wandering wires: frequency of sternal wire abnormalities in patients with sternal dehiscence. *AJR Am J Roentgenol*. 1999;173(3):777–80.
7. Ray JA, Doddi N, Regula D, Williams JA, Melveger A. Polydioxanone (PDS), a novel monofilament synthetic absorbable suture. *Surg Gynecol Obstet*. 1981;153(4):497–507.
8. Keçelgil HT, Kolbaktır F, Akar H, Konuralp C, Demir Z, Demirağ MK. Sternal closure with resorbable synthetic loop suture material in children. *J Pediatr Surg*. 2000;35(9):1309–11.
9. KREITMANN B, Riberi A, METRAS D. Evaluation of an absorbable suture for sternal closure in pediatric cardiac surgery. *J Card Surg*. 1992;7(3):254–6.
10. Hayward RH, Knight WL, Baisden CE, Reiter CG. Sternal dehiscence: early detection by radiography. *J Thorac Cardiovasc Surg*. 1994;108(4):616–9.
11. Bigdelian H, Sedighi M. Evaluation of sternal closure with absorbable polydioxanone sutures in children. *J Cardiovasc Thorac Res*. 2014;6(1):57.
12. Losanoff JE, Basson MD, Gruber SA, Huff H, Hsieh F. Single wire versus double wire loops for median sternotomy closure: experimental biomechanical study using a human cadaveric model. *Ann Thorac Surg*. 2007;84(4):1288–93.
13. Huh J, Bakaeen F, Chu D, Wall Jr MJ. Transverse sternal plating in secondary sternal reconstruction. *J Thorac Cardiovasc Surg*. 2008;136(6):1476–80.
14. Vestergaard R. Sternal Healing Characteristics. Animal and clinical experimental investigation.

- 2013;
15. Claes LE, Heigele CA, Neidlinger-Wilke C, Kaspar D, Seidl W, Margevicius KJ, et al. Effects of mechanical factors on the fracture healing process. *Clin Orthop Relat Res.* 1998;355:S132–47.
 16. El-Ansary D, Waddington G, Adams R. Relationship between pain and upper limb movement in patients with chronic sternal instability following cardiac surgery. *Physiother Theory Pract.* 2007;23(5):273–80.
 17. El-Ansary D, Waddington G, Adams R. Measurement of non-physiological movement in sternal instability by ultrasound. *Ann Thorac Surg.* 2007;83(4):1513–6.
 18. Kurniawan SD. PERBANDINGAN MEKANIK PADA PENUTUPAN STERNUM KAMBING JAWA (*Copra Hircus*) PASCA STERNOTOMI MEDIANA MENGGUNAKAN STAINLESS STEEL WIRE JAHITAN FIGURE OF EIGHT DENGAN POLYDIOXANONE (PDS) JAHITAN SIMPLE CONTINUOUS. Universitas Airlangga; 2016.
 19. Wu LC, Renucci J, Song DH. Rigid-plate fixation for the treatment of sternal nonunion. *J Thorac Cardiovasc Surg.* 2004;128(4):623–4.