

# Intravenous Infiltration and Extravasations: Knowledge among Nurses in Pediatric Hospitals in Khartoum, Sudan

Howaida Mohammed Gasmalla<sup>1</sup>, Mohammed Khalid Hussein<sup>2</sup>, Abdelmoniem Mohamed Hamid<sup>3</sup>,  
Hammad Ali Fadlalmola<sup>4</sup>

<sup>1</sup>Assistant Professor, Faculty of Nursing Sciences, Al Neelain University, Khartoum, Sudan, <sup>2</sup>Assistant Professor, Faculty of Nursing Sciences, Jazan University, Jazan, KSA, <sup>3</sup>Associate Professor, MBBS, MRCPCH (UK), Al Neelain University, Khartoum, Sudan, <sup>4</sup>Associate Professor, Nursing College, Taibah University, KSA

## Abstract

**Introduction:** Infiltration and extravasation are complications of intravenous administration therapies involving unintended leakage of solution into the surrounding tissue. Consequences range from local irritation to amputation. The aim of the study is to study nurses' knowledge about intravenous infiltration and extravasation.

**Method:** Descriptive cross-sectional, hospital-based study. A study was conducted at four pediatric hospitals in Khartoum state. 165 nurses were included using simple random sampling of different working experiences. Data were collected by face-to-face interview questionnaire.

**Results:** Of participants 25.5% defined intravenous infiltration correctly, most participants 76.4% had a poor level of knowledge about identification of infiltration, only 1.8% of participants knew the term extravasation, and most nurses 78.2% had a poor level of knowledge that antibiotics causing extravasation. There was statistically significant with qualification P-value < 0.001

**Conclusion:** The study concluded that nurses had a poor level of knowledge on intravenous infiltration and extravasation. The study recommended education, guideline, and standards for intravenous therapy should be adopted to ensure the best practices.

**Keywords:** *Intravenous infiltration, extravasation, knowledge, nurses, Sudan.*

## Introduction

Intravenous infiltration and extravasation are the most common intravenous therapy complications. Infiltration is the leakage of a non vesicant solution into

the surrounding tissue and extravasation is inadvertent leakage of a vesicant solution into surrounding tissue [1]-[3]. Vesicant refers to any medication or fluid with the potential for causing blisters, severe tissue injury or necrosis [2]. Lack of the knowledge in these vesicants might increase the risk of extravasations [4]. The nurse should routinely assess all vascular access sites for signs and symptoms of infiltration and extravasation based on patient population, type of therapy, type of device, and risk factors [5]. The nurse should determine possible causes of infiltration and extravasation, which include mechanical, pharmacologic, obstructive, and

---

### Corresponding author:

Hammad Ali Fadlalmola<sup>1</sup>, RN, CHN, PhD.  
Associate Professor Taibah University  
Nursing College Department of Community health  
Nursing Saudi Arabia, Madinah Munawara  
E-mail: hazzminno345@gmail.com  
Mobile: 00966504900120

inflammatory factors. It is important for the nurse to be able to recognize the early signs and symptoms of infiltration and extravasation [6]. Extravasation can be caused by physiochemical factors, including pH and osmolarity of some drugs, mechanical factors, occurring either during initial cannula insertion, children may be more susceptible to peripheral intravenous infiltration and extravasation because of developmental and physiological factors, such as communication skills, activity level, and fragile vein [7]. The Infusion Nurses Society (INS) developed an infiltration scale that measured edema in inches and graded numerically from zero to four [3]. A grading scale is recommended for assessing and determining the extent, standardizing the description of the infiltration, documenting the severity of the problem, and evaluating the degree and prevalence of infiltration [8].

It is important to assess the ability of nurses to create positive change in the knowledge and practice [9]-[11]. Number of research regarding intravenous infiltration and extravasation among children from Sudan are limited. It is estimated that 78% of pediatric peripheral intravenous (PIV) lines become infiltrated and 11% of neonatal intensive care unit patients have IV extravasations [3]. The outcome from an infiltrate event can range from edema in an extremity to full-thickness skin loss, muscle, or tendon necrosis, and in some cases, even amputation, these problems not only increased prolonged hospitalization and increased medical costs but also resulted in permanent damage and limitations of physical functions in pediatric patients [12]. This study aimed to assess nurses' knowledge regarding intravenous infiltration and extravasation.

The nursing role is an application of proper techniques which can be used to avoid many cases of complications, such as selecting appropriate intravenous site, checking the catheter before,

during, and after administration of vesicants, avoiding unnecessary coverage of the insertion site, to recognize the early signs and symptoms of infiltration and extravasation, through systematic hourly intravenous site assessments, these various measures help to prevent or limit further damage to the vein and surrounding tissues. It is necessary that the nurses have the knowledge, skills and technical skills that are acquired through training programs and through educational actions [13]-[15].

### **Methodology**

A Descriptive cross-sectional, hospital based-study was conducted at the main pediatric governmental hospitals in Khartoum state (Ahmed Gasim pediatric hospital, Mohammed Elamin Hamid pediatric hospital, Gafaar Ibnouf pediatric referral hospital and ALbluck pediatric hospital) 165 nurses enrolled in the study by using a simple random sampling technique, regard less of qualifications and years of experiences, the sample size was calculated based on total population. Variables under study were, recognition of infiltration and extravasation, pharmacological factors contributing to the risk for extravasation, noncytotoxic medication associated with extravasation and the consequence of IV infiltration and extravasation. Data was gathered through face-to-face interviews questionnaire which composed of two sections: demographic data and knowledge of nurses based on Infusion Therapy Standards of Practice. Knowledge scored according to Likert scale three points, from one to three, 3 being good, 2 for fair and 1 for poor for each question has an answer with more than or equal to three values. Data was analyzed by a computerized program (SPSS) version 20 and presented in the form of simple frequency table and cross table to explore the relationship between variables. P value = 0.05 was considered statistically significant. Ethical approval was obtained from ethical committees and

administrative authorities of hospitals.

## Results

**Table 1. Shows participants demographic characteristics (n=165)**

Demographic	Frequency	Percentage
<b>Age groups</b>		
20-30 years	11	6.70%
31-40 years	92	55.80%
41-50 years	39	23.60%
> 50 years	23	13.90%
<b>Nurses qualifications</b>		
Ordinary nurse	57	34.50%
Diploma	71	43.00%
Bachelor	31	18.80%
Master	6	3.60%
<b>Gender</b>		
Male	30	18.20%
Female	135	81.80%
<b>Years of Experience</b>		
1-5 years	11	6.70%
6-10 years	31	18.80%
11-15 years	86	52.10%
> 15 years	37	22.40%
Total	165	100%

**Table (2): nurses' level of knowledge regarding IV infiltration and extravasation (n=165)**

Variables	knowledge level					
	Poor		Fair		Good	
	N	%	N	%	N	%
Correct definition of IV infiltration	100	60.6	23	13.9	42	25.5
Signs of IV infiltration	126	76.4	17	10.3	22	13.3
Correct definition of extravasation	3	1.8	8	4.8	154	93.3
Signs of extravasation	149	90.3	9	5.5	7	4.2
Pharmacological factors risk for extravasation	153	92.7	10	6.1	2	1.2
Antibiotics associated with extravasation	129	78.2	30	18.2	6	3.6
Electrolyte solutions associated with extravasation	137	83.0	18	10.9	10	6.1
Consequence of IV infiltration and extravasation	150	90.9	11	6.7	4	2.4

**Table 3. The association between qualification and knowledge about definition of extravasation (n=165)**

Qualification		Level of knowledge			Total
		Good	Fair	Poor	
Ordinary nurse	Number	0	0	57	57
	%	0%	0%	100%	100%
Diploma	Number	0	1	70	71
	%	0%	1.4	98.6	100%
BSC	Number	1	3	27	31
	%	3.3%	9.7%	87%	100%
M.Sc.	Number	2	4	0	6
	%	33.4%	66.6%	0%	100%
Total	Number	3	8	154	165
	%	1.8%	4.9%	93.3%	100%

Significant P value 0.001

## Discussion

The study revealed that, quarter (43%) of the nurses carrying a Diploma degree in nursing and only 3.6% had master's degree, probably this because most of MSC holders work at the higher education ministry as lecturers or emigrated. Only (9%) of nurses received a training course about intravenous therapy and its complications, although they have an important role in maintaining IV therapy using their knowledge-about infiltration signs and extravasation, this not similar to study with Lavery he reported that it is important to educate and train nurses associated with IV therapy which is necessary skills for nurses to anticipate problems and take actions to prevent adverse events occurring [16]. Most nurses (76.4%) did not know the signs of infiltration, this disagrees with the study done by Dougherty [12]. Regarding the definition of extravasation only (1.8%) of nurses defined extravasation correctly, most of the nurses (93.3%) did not know the term extravasation this disagree to study by Sisan 54% of nurses knew the correct definition for extravasation and infiltration [17]. The nurses failed to define the term extravasation; this finding may be related to lack of education. (4.2%) of nurses only had a good level about the information that redness followed by blistering and necrosis signs of extravasation. In Sudan many of children with black skin color, which is difficult to recognize extravasation early, this is supported by Treadwell that children and neonates with darker skin are more likely to suffer from extravasation [18]. Moreover (83.0%) nurses had a poor level of knowledge about electrolytes associated with extravasation. Doellman reported that Calcium Chloride leads to ischemia [19]. There was an inconsistency of nurses' knowledge regarding non cytotoxic medications associated with extravasation ,their good level of knowledge about antibiotics causing extravasation was (3.6%), this finding due to lack of in service workshops and

training courses , in addition poor level of knowledge (90.9%) of the nurses about consequences of infiltration and extravasation this not in context to study done by Park reported that the injury associated with extravasation in worst situations may require amputation, so the nurse must be vigilant for early signs of infiltration and extravasation [20].The current study showed differences in level of knowledge to nurses qualification p-value < 0.001.

## Conclusion

The study revealed that IV infiltration and extravasation knowledge among nurses at pediatric hospitals in Khartoum State was poor, and poor knowledge is significantly associated with the qualification of nurses. The study recommends that education, guidelines, and standards for intravenous therapy should be adopted to ensure the best practice.

**Conflicts of Interest:** The authors declare that there are no conflicts of interest

**Financial Support:** There was no support for this research from any funding source, commercial or non-profit.

**Ethical Clearance:** Ethical approval was obtained from ethical committees and administrative authorities of hospitals.

## References

1. Dychter SS, Gold DA, Carson D, et al. Intravenous therapy: a review of complications and economic considerations of peripheral access. *Journal of Infusion Nursing* 2012; 35: 84–91.
2. Hadaway L. Infiltration and extravasation. *AJN The American Journal of Nursing* 2007; 107: 64–72.
3. Doellman D, Hadaway L, Bowe-Geddes LA, et al. Infiltration and extravasation: update on prevention and management. *Journal of Infusion*

- Nursing* 2009; 32: 203–211.
4. Thomas J. Standard practice and evolving trends in pediatric intravenous access. *Air medical journal* 2007; 26: 8–11.
  5. Infusions I. Site Care and Maintenance.
  6. Lyons MG, Kasker J. Outcomes of a continuing education course on intravenous catheter insertion for experienced registered nurses. *The Journal of Continuing Education in Nursing* 2012; 43: 177–181.
  7. Alexander M. Infusion Standards: A Document Without Borders. *Journal of Infusion Nursing* 2016; 39: 181–182.
  8. Abdelaziz R Ben, Hafsi H, Hajji H, et al. Full title: peripheral venous catheter complications in children: predisposing factors in a multicenter prospective cohort study. *BMC pediatrics* 2017; 17: 1–11.
  9. Majid S, Foo S, Luyt B, et al. Adopting evidence-based practice in clinical decision making: nurses' perceptions, knowledge, and barriers. *Journal of the Medical Library Association: JMLA* 2011; 99: 229.
  10. Malik G, McKenna L, Plummer V. Perceived knowledge, skills, attitude and contextual factors affecting evidence-based practice among nurse educators, clinical coaches and nurse specialists. *International journal of nursing practice* 2015; 21: 46–57.
  11. Ibrahim MM, Fadlalmola HA. Effects of Nurse 's Work Environment and Practice on Patient 's Safety. 2020; 15: 345–354.
  12. Dougherty L. IV therapy: recognizing the differences between infiltration and extravasation. *British Journal of Nursing* 2008; 17: 896–901.
  13. Tofani BF, Rineair SA, Gosdin CH, et al. Quality improvement project to reduce infiltration and extravasation events in a pediatric hospital. *Journal of pediatric nursing* 2012; 27: 682–689.
  14. Clifton-Koeppel R. Wound care after peripheral intravenous extravasation: What is the evidence? *Newborn and Infant Nursing Reviews* 2006; 6: 202–211.
  15. Verity R, Wiseman T, Ream E, et al. Exploring the work of nurses who administer chemotherapy. *European Journal of Oncology Nursing* 2008; 12: 244–252.
  16. Lavery I. Infection control in IV therapy: a review of the chain of infection. *British journal of nursing* 2010; 19: S6–S14.
  17. McCullen KL, Pieper B. A retrospective chart review of risk factors for extravasation among neonates receiving peripheral intravascular fluids. *Journal of Wound Ostomy & Continence Nursing* 2006; 33: 133–139.
  18. Sauerland C, Engelking C, Wickham R, et al. Vesicant extravasation part I: mechanisms, pathogenesis, and nursing care to reduce risk. In: *Oncology nursing forum*. 2006.
  19. Beaulieu MJ. Hyaluronidase for extravasation management. *Neonatal network* 2012; 31: 413–419.
  20. Park SM, Jeong IS, Kim KL, et al. The effect of intravenous infiltration management program for hospitalized children. *Journal of pediatric nursing* 2016; 31: 172–178.